arXiv:1207.5842v1 [math.DS] 24 Jul 2012

QUANTIZATION DIMENSION FOR GIBBS-LIKE MEASURES ON COOKIE-CUTTER SETS

MRINAL KANTI ROYCHOWDHURY

Abstract. In this paper using Banach limit we have determined a Gibbs-like measure μ_h supported by a cookie-cutter set E which is generated by a single cookie-cutter mapping f. For such a measure μ_h and a given $r \in (0, +\infty)$ we have determined the quantization dimension function $D_r := D_r(\mu_h)$, and established its functional relationship with the temperature function of the thermodynamic formalism. The temperature function is commonly used to perform the multifractal analysis, in our context of the measure μ_h.

1. Introduction

Quantization dimension is one of the most important objects in the quantization problem, which has a deep background in information theory and engineering technology (cf. [BW, GG, GN, Z]). It characterizes in a natural way the asymptotic property of the error when approximating a given probability measure by a discrete probability measure of finite support in the sense of L_r-metrics. Given a Borel probability measure μ on $\mathbb{R}^d$, a number $r \in (0, +\infty)$ and a natural number $n \in \mathbb{N}$, the nth *quantization error* of order r for μ, is defined by

$$V_{n,r}(\mu) := \inf\{\int d(x, \alpha)^r d\mu(x) : \alpha \subset \mathbb{R}^d, \text{ card}(\alpha) \leq n\},$$

where $d(x, \alpha)$ denotes the distance from the point x to the set α with respect to a given norm $\| \cdot \|$ on $\mathbb{R}^d$. Note that if $\int \|x\|^r d\mu(x) < \infty$ then there is some set α for which the infimum is achieved (cf. [GL1]). The set α for which the infimum is achieved is called an optimal set of n-means or n-optimal set of order r for $0 < r < +\infty$. The upper and lower quantization dimension of order r for μ is defined to be

$$\overline{D}_r(\mu) := \limsup_{n\to\infty} \frac{r \log n}{-\log V_{n,r}(\mu)}; \ \underline{D}_r(\mu) := \liminf_{n\to\infty} \frac{r \log n}{-\log V_{n,r}(\mu)}.$$

If $\overline{D}_r(\mu)$ and $\underline{D}_r(\mu)$ coincide, we call the common value the *quantization dimension of order* r for the probability measure μ, and is denoted by $D_r := D_r(\mu)$. One sees that the quantization dimension is actually a function $r \mapsto D_r$ which measures the asymptotic rate at which $V_{n,r}$ goes to zero. If D_r exists, then one can write

$$\log V_{n,r} \sim \log(\frac{1}{n})^{r/D_r}.$$

For probabilities with non-vanishing absolutely continuous part the numbers D_r are all equal to the dimension d of the underlying space, but for singular probabilities the family $(D_r)_{r>0}$ gives an interesting description of their geometric (multifractal) structures.

Let $S_1, S_2, \cdots, S_N$ be contractive similitudes from $\mathbb{R}^d$ into itself, where $N \geq 2$ is a positive integer. Let s_i be the contraction ratio of S_i for all $1 \leq i \leq N$. Then for a given probability

---

2000 *Mathematics Subject Classification.* 60Exx, 28A80, 94A34.

Key words and phrases. Gibbs like measure, quantization dimension, topological pressure, temperature function.

vector $(p_1, p_2, \cdots, p_N)$ there exists a unique Borel probability measure μ (cf. [H]) satisfying the condition

$$\mu = \sum_{j=1}^{N} p_j \mu \circ S_j^{-1}.$$

Let the iterated function system $\{S_1, S_2, \cdots, S_N\}$ satisfy the open set condition: there exists a bounded nonempty open set $U \subset \mathbb{R}^d$ such that $\bigcup_{j=1}^{N} S_j(U) \subset U$ and $S_i(U) \bigcap S_j(U) = \emptyset$ for $1 \leq i \neq j \leq N$. The iterated function system satisfies the strong open set condition if U can be chosen such that $U \cap J \neq \emptyset$, where J is the limit set of the iterated function system. Under the open set condition, Graf and Luschgy showed that the quantization dimension function $D_r := D_r(\mu)$ of the probability measure μ exists, and satisfies the following relation (cf. [GL1, GL2]):

$$\sum_{j=1}^{N} (p_j s_j^r)^{\frac{D_r}{r+D_r}} = 1.$$

In fact, they proved a stronger result namely that the quantization dimension D_r also satisfies (cf. [GL3]):

$$0 < \liminf_{n\to\infty} n V_{n,r}^{\frac{D_r}{r}}(\mu) \leq \limsup n V_{n,r}^{\frac{D_r}{r}}(\mu) < +\infty. \tag{1}$$

Under the open set condition, Lindsay and Mauldin (cf. [LM]) determined the quantization dimension for an F-conformal measure m associated with a conformal iterated function system determined by finitely many conformal mappings. They established a relationship between the quantization dimension and the temperature function of the thermodynamic formalism arising in multifractal analysis. Later, quantization dimension and its relationship with the temperature function were determined for some other fractal probability measures, for example one could see [R1, R2, WD].

In this paper, using Banach limit we have defined a Gibbs-like measure μ_h supported by a cookie-cutter set E, where E is the limit set generated by a cooki-cutter mapping f and $h := \dim_H(E)$ is the Hausdorff dimension of the set E (cf. [F1]). For this measure μ_h we have shown that for each $r \in (0, +\infty)$ there exists a unique $\kappa_r \in (0, +\infty)$ such that

$$\lim_{k\to\infty} \frac{1}{k} \log \sum_{\sigma\in\Omega_k} (\mu_h(J_\sigma) \|\varphi_\sigma'\|^r)^{\frac{\kappa_r}{r+\kappa_r}} = 0, \tag{2}$$

and the above κ_r is the quantization dimension for the probability measure μ_h. It is known that, the singularity exponent $\beta(q)$ (also known as the temperature function) satisfies the usual equation

$$\lim_{k\to\infty} \frac{1}{k} \log \sum_{\sigma\in\Omega_k} (\mu_h(J_\sigma))^q \|\varphi_\sigma'\|^{\beta(q)} = 0, \tag{3}$$

and that the spectrum $f(\alpha)$ is the Legendre transform of $\beta(q)$. Comparing (2) and (3), we see that if $q_r = \frac{D_r}{r+D_r}$, then $\beta(q_r) = rq_r$, that is, the quantization dimension function for the probability measure μ_h has a relationship with the temperature function of the thermodynamic formalism arising in multifractal analysis (for thermodynamic formalism, multifractal analysis and the Legendre transform one could see [F1]).

2. Basic definitions, lemmas and propositions

In this paper, $\mathbb{R}^d$ denotes the d-dimensional Euclidean space equipped with a metric d. Let us write

$$V_{n,r}(\mu) := \inf\{\int d(x,\alpha)^r d\mu(x) : \alpha \subset \mathbb{R}^d,\ \text{card}(\alpha) \leq n\},$$
$$u_{n,r}(\mu) := \inf\{\int d(x,\alpha \cup U^c)^r d\mu(x) : \alpha \subset \mathbb{R}^d,\ \text{card}(\alpha) \leq n\},$$

where U is a set which comes from the open set condition and U^c denotes the complement of U. We see that

$$u_{n,r}^{1/r} \leq V_{n,r}^{1/r} := e_{n,r}.$$

We call sets $\alpha_n \subset \mathbb{R}^d$, for which the above infimums are achieved, n-optimal sets for $e_{n,r}, V_{n,r}$ or $u_{n,r}$ respectively. As stated above, Graf and Luschgy have shown that n-optimal sets exist when $\int \|x\|^r d\mu(x) < \infty$.

2.1. **Cookie-cutter set:** A mapping f is called a cookie-cutter, if there exists a finite collection of disjoint closed intervals $J_1, J_2, \cdots, J_N \subset J = [0,1]$, such that

$(C1)$ f is defined in a neighborhood of each J_j, $1 \leq j \leq N$, the restriction of f to each initial interval J_j is 1-1 and onto, the corresponding branch inverse is denoted by $\varphi_j := (f|_{J_j})^{-1} : J \to J_j$;

$(C2)$ f is differentiable with Hölder continuous derivative f', i.e., there exist constants $c > 0$ and $\gamma \in (0,1]$ such that for $x, y \in J_j$, $1 \leq j \leq N$,

$$|f'(x) - f'(y)| \leq c|x-y|^\gamma;$$

$(C3)$ f is boundedly expanding in the sense that there exist constants b and B

$$1 < b := \inf_x\{|f'(x)|\} \leq \sup_x\{|f'(x)|\} := B < +\infty.$$

$[\bigcup_{j=1}^N J_j; c, \gamma, b, B]$ is called the defining data of the cookie-cutter mapping f. Let Ω_0 be the empty set. For $n \geq 1$, define

$$\Omega_n = \{1,2,\cdots,N\}^n,\ \Omega_\infty = \{1,2,\cdots,N\}^{\mathbb{N}} \text{ and } \Omega = \bigcup_{k=0}^\infty \Omega_k.$$

Elements of Ω are called words. For any $\sigma \in \Omega$ if $\sigma = (\sigma_1, \sigma_2, \cdots, \sigma_n) \in \Omega_n$, we write $\sigma^- = (\sigma_1, \sigma_2, \cdots, \sigma_{n-1})$ to denote the word obtained by deleting the last letter of σ, $|\sigma| = n$ to denote the length of σ, and $\sigma|_k := (\sigma_1, \sigma_2, \cdots, \sigma_k)$, $k \leq n$, to denote the truncation of σ to the length k. For any two words $\sigma = (\sigma_1, \sigma_2, \cdots, \sigma_k)$ and $\tau = (\tau_1, \tau_2, \cdots, \tau_m)$, we write $\sigma\tau = \sigma * \tau = (\sigma_1, \cdots, \sigma_k, \tau_1, \cdots, \tau_m)$ to denote the juxtaposition of $\sigma, \tau \in \Omega$. A word of length zero is called the empty word and is denoted by $\emptyset$. For $\sigma \in \Omega$ and $\tau \in \Omega \cup \Omega_\infty$ we say τ is an extension of σ, written as $\sigma \prec \tau$, if $\tau|_{|\sigma|} = \sigma$. For $\sigma \in \Omega_k$, the cylinder set $C(\sigma)$ is defined as $C(\sigma) = \{\tau \in \Omega_\infty : \tau|_k = \sigma\}$. For $\sigma = (\sigma_1, \sigma_2, \cdots, \sigma_n) \in \Omega_n$, let us write $\varphi_\sigma = \varphi_{\sigma_1} \circ \cdots \circ \varphi_{\sigma_n}$, and define the rank-$n$ basic interval corresponding to σ by

$$J_\sigma = J_{(\sigma_1,\sigma_2,\cdots,\sigma_n)} = \varphi_\sigma(J),$$

where $1 \leq \sigma_k \leq N$, $1 \leq k \leq n$. If $\sigma = \emptyset$, then we identify $\varphi_\emptyset$ to be the identity mapping on J and write $J_\emptyset = J$. By $|J_\sigma|$ we mean the diameter of the set J_σ for all $\sigma \in \Omega$. It is easy to see that the set of basic intervals $\{J_\sigma : \sigma \in \Omega\}$ has the following net properties:

(i) $J_{\sigma * j} \subset J_\sigma$ for each $\sigma \in \Omega_n$ and $1 \leq j \leq N$ for all $n \geq 1$;

(ii) $J_\sigma \bigcap J_\tau = \emptyset$, if $\sigma, \tau \in \Omega_n$ for all $n \geq 1$ and $\sigma \neq \tau$.

Since φ_j is a branch inverse of f, where $1 \leq j \leq N$, for all $x \in J$, we have $f(\varphi_j(x)) = x$, and so $|f'(\varphi_j(x))| \cdot |\varphi_j'(x)| = 1$, which yields

$$B^{-1} \leq \left|\varphi_j'(x)\right| \leq b^{-1}. \tag{4}$$

Choose x, y to be the end points of J, and then $\varphi_\sigma(x), \varphi_\sigma(y)$ are the end points of J_σ for each $\sigma \in \Omega$, and so by mean value theorem,

$$|J_\sigma| = |\varphi_\sigma(x) - \varphi_\sigma(y)| = |\varphi_\sigma'(w)||x - y| = |\varphi_\sigma'(w)|,$$

for some $w \in J_\sigma$. Thus $B^{-n} \leq |J_\sigma| \leq b^{-n}$ for any $\sigma \in \Omega_n$, and thus the diameter $|J_\sigma| \to 0$ as $|\sigma| \to \infty$. Since given $\sigma = (\sigma_i)_{i=1}^\infty \in \Omega_\infty$ the diameters of the compact sets $J_{\sigma|_k}$, $k \geq 1$, converge to zero and since they form a descending family, the set

$$\bigcap_{k=0}^\infty J_{\sigma|_k}$$

is a singleton and therefore, if we denote its element by $\pi(\sigma)$, this defines the coding map $\pi : \Omega_\infty \to J$. The main object of our interest is the limit set

$$E := \pi(\Omega_\infty) = \bigcup_{\sigma\in\Omega_\infty} \bigcap_{k=0}^\infty J_{\sigma|_k}.$$

Moreover, $\pi(C(\sigma)) = E \cap J_\sigma$ for $\sigma \in \Omega$. With the net properties it follows that E is a perfect, nowhere dense and totally disconnected subset of J. The set E is called the *cookie-cutter* set.

Let ℓ^∞ be the set of all bounded sequences $x = (x_n)_{n\in\mathbb{N}}$ of real or complex numbers, which form a vector space with respect to point-wise addition and multiplication by a scalar. It is equipped with the norm $\|x\| = \sup_n |x_n|$. The normed space ℓ^∞ is complete with respect to the metric $\|x - y\|$, and so it forms a Banach space. By the Hahn-Banach theorem (cf. [Y, p. 102-104]), there exists a linear functional $L : \ell^\infty \to \mathbb{R}$ for which

(i) L is linear;
(ii) $L((x_n)_{n\in\mathbb{N}}) = L((x_{n+1})_{n\in\mathbb{N}})$;
(iii) $\liminf_{n\to\infty}(x_n) \leq L((x_n)_{n\in\mathbb{N}}) \leq \limsup_{n\to\infty}(x_n)$.

The functional L, defined above, is called a Banach limit. The use of the Banach limit is rather a standard tool in producing an invariant measure from a given measure.

Let us now prove the following lemma.

Lemma 2.2. ***(Bounded variation principle)*** *There exists a constant $1 < \xi < +\infty$ such that for each $\sigma \in \Omega_n$, and $x, y \in J_\sigma$, we have*

$$\xi^{-1} \leq \frac{|(f^n)'(x)|}{|(f^n)'(y)|} \leq \xi,$$

where $f^n = f \circ f \circ \cdots \circ f$ represents the n-fold composition of f with itself.

Proof. Note that for each $k \leq n$ and $\sigma = (\sigma_1, \sigma_2, \cdots, \sigma_n) \in \Omega_n$, f^{k-1} maps J_σ diffeomorphically to the set $\varphi_{\sigma_k} \circ \varphi_{\sigma_{k+1}} \circ \cdots \circ \varphi_{\sigma_n}(J)$, and so

$$\left|f^{k-1}(x) - f^{k-1}(y)\right| \leq \operatorname{diam}\left(\varphi_{\sigma_k} \circ \varphi_{\sigma_{k+1}} \circ \cdots \circ \varphi_{\sigma_n}(J)\right) = |\varphi_{\sigma_k} \circ \varphi_{\sigma_{k+1}} \circ \cdots \circ \varphi_{\sigma_n}(J)|.$$

By mean value theorem,

$$\begin{aligned}
&|\varphi_k \circ \varphi_{\sigma_{k+1}} \circ \cdots \circ \varphi_{\sigma_n}(J)| \\
&= \sup_{x,y\in J} \left|\varphi_{\sigma_k}\left(\varphi_{\sigma_{k+1}} \circ \cdots \circ \varphi_{n,\sigma_n}(x)\right) - \varphi_{\sigma_k}\left(\varphi_{\sigma_{k+1}} \circ \cdots \circ \varphi_{\sigma_n}(y)\right)\right| \\
&\leq b^{-1}\left|\varphi_{\sigma_{k+1}} \circ \cdots \circ \varphi_{\sigma_n}(J)\right|.
\end{aligned}$$

Thus proceeding inductively,

$$\left|\varphi_{\sigma_k}\circ\varphi_{\sigma_{k+1}}\circ\cdots\circ\varphi_{\sigma_n}(J)\right|\leq b^{-(n-k+1)}.$$

Then, Hölder continuity of f' gives

$$\left|f'(f^{k-1}(x))-f'(f^{k-1}(y))\right|\leq c\left|f^{k-1}(x)-f^{k-1}(y)\right|^{\gamma}\leq cb^{-(n-k+1)\gamma},$$

and so by mean value theorem and the assumption $|f'|>1$, we have

$$\begin{aligned}&\Big|\log|f'(f^{k-1}(x))|-\log|f'(f^{k-1}(y))|\Big|\\ &\leq\Big||f'(f^{k-1}(x))|-|f'(f^{k-1}(y))|\Big|\\ &\leq cb^{-(n-k+1)\gamma}.\end{aligned}$$

Therefore, by the above inequality and the chain rule,

$$\begin{aligned}&\Big|\log|(f^n)'(x)|-\log|(f^n)'(y)|\Big|\\ &=\Big|\sum_{k=1}^{n}\log|f'(f^{k-1}(x))|-\sum_{k=1}^{n}\log|f'(f^{k-1}(y))|\Big|\\ &\leq\sum_{k=1}^{n}\Big|\log|f'(f^{k-1}(x))|-\log|f'(f^{k-1}(y))|\Big|\\ &\leq\sum_{k=1}^{n}cb^{-(n-k+1)\gamma}\leq\frac{cb^{-\gamma}}{1-b^{-\gamma}}.\end{aligned}$$

Take $\xi=\exp\left\{\frac{c}{b^{\gamma}-1}\right\}$. Since $\frac{c}{b^{\gamma}-1}>0$, we have $1<\xi<+\infty$, and thus the proposition follows. ☐

Let us now prove the following proposition.

Proposition 2.3. . (**Bounded distortion principle**) For any $n\geq 1$, $\sigma\in\Omega_n$, $x\in J_\sigma$, we have

$$\xi^{-1}\leq|(f^n)'(x)|\cdot|J_\sigma|\leq\xi.$$

Moreover, for each $1\leq j\leq N$, we get $|J_{\sigma*j}|\geq\xi^{-2}B^{-1}|J_\sigma|$, where ξ is the constant of Lemma 2.2.

Proof. Note that for $\sigma\in\Omega_n$, $f^n:J_\sigma\to J$ is a differentiable bijection. So by mean value theorem, if $y,z\in J_\sigma$, there exists $w\in J_\sigma$ such that

$$f^n(y)-f^n(z)=(f^n)'(w)(y-z).$$

Choose y,z to be the end points of J_σ, and then $f^n(y),f^n(z)$ are the end points of J, and so

$$|J|=|(f^n)'(w)|\cdot|J_\sigma|,\text{ i.e. }|(f^n)'(w)|\cdot|J_\sigma|=1.$$

Hence, using bounded variation principle, we have

$$\xi^{-1}\leq|(f^n)'(x)|\cdot|J_\sigma|\leq\xi \tag{5}$$

for all $x\in J_\sigma$. Now let $1\leq j\leq N$ and $x\in J_{\sigma*j}$. Then using (5), we have

$$\xi^{-1}\leq|(f^{n+1})'(x)|\cdot|J_{\sigma*j}|=|(f'(f^n(x))|\cdot|(f^n)'(x)|\cdot|J_{\sigma*j}|\leq B|(f^n)'(x)|\cdot|J_{\sigma*j}|.$$

Since $J_{\sigma*j}\subseteq J_\sigma$, we have $x\in J_\sigma$. Hence using (5) again, we have

$$|J_{\sigma*j}|\geq\xi^{-2}B^{-1}|J_\sigma|.$$

Thus the proof of the proposition is yielded. ☐

Proposition 2.4. For any $n \geq 1$, let $\sigma \in \Omega_n$, and $x, y \in J$. Let ξ be the constant of Lemma 2.2. Then,

$$\xi^{-1}|\varphi'_\sigma(y)| \leq |\varphi'_\sigma(x)| \leq \xi|\varphi'_\sigma(y)|.$$

Proof. For $\sigma \in \Omega_n$ and $x \in J$, we know $f^n(\varphi_\sigma(x)) = x$. Thus

$$|(f^n)'(\varphi_\sigma(x))| \cdot |\varphi'_\sigma(x)| = 1.$$

Again for all $x \in J$, $\varphi_\sigma(x) \in J_\sigma$. Hence, Lemma 2.2 yields

$$\xi^{-1}|\varphi'_\sigma(y)| \leq |\varphi'_\sigma(x)| \leq \xi|\varphi'_\sigma(y)|,$$

and thus the proposition is obtained. ☐

Let us now prove the following two lemmas.

Lemma 2.5. *Let $\sigma, \tau \in \Omega$. Then*

$$\xi^{-1}\|\varphi'_\sigma\|\|\varphi'_\tau\| \leq \|\varphi'_{\sigma\tau}\| \leq \|\varphi'_\sigma\|\|\varphi'_\tau\|,$$

ξ is the constant of Lemma 2.2

Proof. For any $x \in J$, we have $|\varphi'_{\sigma\tau}(x)| = |\varphi'_\sigma(\varphi_\tau(x))| \cdot |\varphi'_\tau(x)|$, and thus by Proposition 2.4, for any $y \in J$,

$$\xi^{-1}|\varphi'_\sigma(y)|\cdot|\varphi'_\tau(x)| \leq |\varphi'_\sigma(\varphi_\tau(x))|\cdot|\varphi'_\tau(x)| = |\varphi_{\sigma\tau}(x)| \leq |\varphi'_\sigma(\varphi_\tau(x))|\cdot|\varphi_\tau(x)| \leq \xi|\varphi'_\sigma(y)|\cdot|\varphi'_\tau(x)|,$$

and thus the lemma follows. ☐

Lemma 2.6. *Let $\sigma \in \Omega$ and $x \in J$. Then*

$$\xi^{-1}|J_\sigma| \leq |\varphi'_\sigma(x)| \leq \xi|J_\sigma|,$$

where ξ is the constant of Lemma 2.2.

Proof. Let $x \in J$, and then $\varphi_\sigma(x) \in J_\sigma$ for $\sigma \in \Omega_n$, $n \geq 1$. We know $f^n(\varphi_\sigma(x)) = x$, and so $|(f^n)'(\varphi_\sigma(x))| \cdot |\varphi'_\sigma(x)| = 1$. Now use Proposition 2.3 to obtain the lemma. ☐

Let us now prove the following lemma.

Lemma 2.7. *Let $\sigma, \tau \in \Omega$. Then*

$$\xi^{-3}|J_\sigma||J_\tau| \leq |J_{\sigma\tau}| \leq \xi^3|J_\sigma||J_\tau|,$$

where ξ is the constant of Lemma 2.2.

Proof. For $\sigma, \tau \in \Omega$, we have $|\varphi'_{\sigma\tau}(x)| = |\varphi'_\sigma(y)||\varphi'_\tau(x)|$ where $y = \varphi_\tau(x)$, and $x \in J$. Again Proposition 2.4 gives that for any $x, y \in J$

$$\xi^{-1}|\varphi'_\sigma(y)| \leq |\varphi'_\sigma(x)| \leq \xi|\varphi'_\sigma(y)|.$$

Hence, Lemma 2.6 implies

$$\xi^{-3}|J_\sigma||J_\tau| \leq \xi^{-1}|\varphi'_\sigma(y)||\varphi'_\tau(x)| = \xi^{-1}|\varphi'_{\sigma\tau}(x)| \leq |J_{\sigma\tau}| \leq \xi|\varphi'_{\sigma\tau}(x)| \leq \xi^3|J_\sigma||J_\tau|,$$

and thus the lemma is obtained. ☐

By Lemma 2.5 and the standard theory of sub-additive sequences, the function $Q(t)$ given by

$$Q(t) = \lim_{k\to\infty} \frac{1}{k} \log \sum_{\sigma\in\Omega_k} \|\varphi'_\sigma\|^t,$$

for any real t exists. It is easy to see that the function $Q(t)$ is strictly decreasing convex and hence continuous in t.

Lemma 2.8. *There exists a unique $h \in (0, +\infty)$ such that $Q(h) = 0$.*

Proof. Since the function $Q(t)$ is strictly decreasing and continuous on $\mathbb{R}$, there exists a unique $h \in \mathbb{R}$ such that $Q(h) = 0$. Note that

$$Q(0) = \lim_{k\to\infty} \frac{1}{k} \log \sum_{\sigma\in\Omega_k} 1 \geq \lim_{k\to\infty} \frac{1}{k} \log N^k = \log N \geq \log 2 > 0.$$

In order to conclude the proof it therefore suffices to show that $\lim_{t\to+\infty} Q(t) = -\infty$. If $t > 0$,

$$Q(t) = \lim_{k\to\infty} \frac{1}{k} \log \sum_{\sigma\in\Omega_k} \|\varphi'_\sigma\|^t \leq \lim_{k\to\infty} \frac{1}{k} \log \sum_{\sigma\in\Omega_k} b^{-kt} = \lim_{k\to\infty} \frac{1}{k} \log N^k - t\log b = \log N - t\log b.$$

Since $b > 1$, it follows that $\lim_{t\to+\infty} Q(t) = -\infty$, and hence the lemma follows. □

Note 2.9. Lemma 2.6 implies $\xi^{-1}|J_\sigma| \leq \sup_{x\in J} |\varphi'_\sigma(x)| = \|\varphi'_\sigma\| \leq \xi|J_\sigma|$, and so the topological pressure $Q(t)$ can be written as follows:

$$Q(t) = \lim_{k\to\infty} \frac{1}{k} \log \sum_{\sigma\in\Omega_k} |J_\sigma|^t.$$

The unique $h \in (0, +\infty)$ given by Lemma 2.8 is the Hausdorff dimension $\dim_H(E)$ of the cookie-cutter set E (cf. [F1]).

Let us now prove the following proposition, which plays a vital role in the paper.

Proposition 2.10. *Let $h \in (0, +\infty)$ be unique such that $Q(h) = 0$, and let s_* and s^* be any two arbitrary real numbers with $0 < s_* < h < s^*$. Then for all $n \geq 1$,*

$$\xi^{-3s_*} < \sum_{\sigma\in\Omega_n} |J_\sigma|^{s_*} \text{ and } \sum_{\sigma\in\Omega_n} |J_\sigma|^{s^*} < \xi^{3s^*},$$

where ξ is the constant of Lemma 2.2.

Proof. Let $s_* < h$. As the pressure function $Q(t)$ is strictly decreasing, $Q(s_*) > Q(h) = 0$. Then for any positive integer n, by Lemma 2.7, we have

$$0 < Q(s_*) = \lim_{p\to\infty} \frac{1}{np} \log \sum_{\omega\in\Omega_{np}} |J_\omega|^{s_*} \leq \lim_{p\to\infty} \frac{1}{np} \log \xi^{3(p-1)s_*} \left(\sum_{\sigma\in\Omega_n} |J_\sigma|^{s_*} \right)^p,$$

which implies

$$0 < \frac{1}{n} \log \left(\xi^{3s_*} \sum_{\sigma\in\Omega_n} |J_\sigma|^{s_*} \right) \text{ and so } \sum_{\sigma\in\Omega_n} |J_\sigma|^{s_*} > \xi^{-3s_*}.$$

Now if $h < s^*$, then $Q(s^*) < 0$ as $Q(t)$ is strictly decreasing. Then for any positive integer n, by Lemma 2.7, we have

$$0 > Q(s^*) = \lim_{p\to\infty} \frac{1}{np} \log \sum_{\omega\in\Omega_{np}} |J_\omega|^{s^*} \geq \lim_{p\to\infty} \frac{1}{np} \log \xi^{-3(p-1)s^*} \left(\sum_{\sigma\in\Omega_n} |J_\sigma|^{s^*} \right)^p,$$

which implies

$$0 > \frac{1}{n} \log \left(\xi^{-3s^*} \sum_{\sigma\in\Omega_n} |J_\sigma|^{s^*} \right) \text{ and so } \sum_{\sigma\in\Omega_n} |J_\sigma|^{s^*} < \xi^{3s^*}.$$

Thus the proposition is obtained. □

Corollary 2.11. *Since s_* and s^* be any two arbitrary real numbers with $0 < s_* < h < s^*$, from the above proposition it follows that for all $n \geq 1$,*

$$\xi^{-3h} \leq \sum_{\sigma\in\Omega_n} |J_\sigma|^h \leq \xi^{3h}.$$

Let us now prove the following proposition.

Proposition 2.12. (Gibbs-like measure) Let $h \in (0, +\infty)$ be such that $Q(h) = 0$. Then there exists a constant $\eta > 1$ and a probability measure μ_h supported by E such that for any $\sigma \in \Omega$,

$$\eta^{-1}|J_\sigma|^h \leq \mu_h(J_\sigma) \leq \eta|J_\sigma|^h.$$

Proof. For $\sigma \in \Omega$, $n \geq 1$, define

$$\nu_n(C(\sigma)) = \frac{\sum_{\tau\in\Omega_n}(\text{diam}J_{\sigma\tau})^h}{\sum_{\tau\in\Omega_{|\sigma|+n}}(\text{diam}J_\tau)^h}.$$

Then using Lemma 2.7 and Corollary 2.11, we have

$$\nu_n(C(\sigma)) \leq \frac{\xi^{3h}(\text{diam}J_\sigma)^h\sum_{\tau\in\Omega_n}(\text{diam}J_\tau)^h}{\sum_{\tau\in\Omega_{|\sigma|+n}}(\text{diam}J_\tau)^h} \leq \xi^{9h}(\text{diam}J_\sigma)^h,$$

and similarly, $\nu_n(C(\sigma)) \geq \xi^{-9h}(\text{diam}J_\sigma)^h$. Thus for a given $\sigma \in \Omega$, $\{\nu_n(C(\sigma))\}_{n=1}^\infty$ is a bounded sequence of real numbers, and so Banach limit, denoted by Lim, is defined. For $\sigma \in \Omega$, let

$$\nu(C(\sigma)) = \text{Lim}_{n\to\infty}\nu_n(C(\sigma)).$$

Then

$$\sum_{j=1}^N \nu(C(\sigma j)) = \text{Lim}_{n\to\infty}\sum_{j=1}^N \frac{\sum_{\tau\in\Omega_n}(\text{diam}J_{\sigma j\tau})^h}{\sum_{\tau\in\Omega_{|\sigma|+1+n}}(\text{diam}J_\tau)^h} = \text{Lim}_{n\to\infty}\frac{\sum_{\tau\in\Omega_{n+1}}(\text{diam}J_{\sigma\tau})^h}{\sum_{\tau\in\Omega_{|\sigma|+n+1}}(\text{diam}J_\tau)^h},$$

and so

$$\sum_{j=1}^N \nu(C(\sigma j)) = \text{Lim}_{n\to\infty}\nu_{n+1}(C(\sigma)) = \text{Lim}_{n\to\infty}\nu_n(C(\sigma)) = \nu(C(\sigma)).$$

Thus by Kolmogorov's extension theorem, ν can be extended to a unique Borel probability measure γ on Ω_∞. Let μ_h be the image measure of γ under the coding map π, i.e., $\mu_h = \gamma\circ\pi^{-1}$. Then μ_h is a unique Borel probability measure supported by E. Moreover, for any $\sigma \in \Omega$,

$$\mu_h(J_\sigma) = \gamma(C(\sigma)) = \text{Lim}_{n\to\infty}\nu_n(C(\sigma)) \leq \text{Lim}_{n\to\infty}\xi^{9h}(\text{diam}J_\sigma)^h = \xi^{9h}(\text{diam}J_\sigma)^h,$$

and similarly,

$$\mu_h(J_\sigma) \geq \xi^{-9h}(\text{diam}J_\sigma)^h.$$

Write $\eta = \xi^{9h}$, and then $\eta > 1$, and thus the proof of the proposition is complete. ☐

For the above measure μ_h, known as *Gibbs-like measure*, we will determine the quantization dimension function and its functional relationship with the temperature function of the thermodynamic formalism.

Let us now prove the following lemma.

Lemma 2.13. *Let μ_h be the Gibbs-like measure as defined in Proposition 2.12. Then there exists a constant $L > 1$ such that for $\sigma, \tau \in \Omega$,*

$$L^{-1}\mu_h(J_\sigma)\mu_h(J_\tau) \leq \mu_h(J_{\sigma\tau}) \leq L\mu_h(J_\sigma)\mu_h(J_\tau).$$

Proof. Let $\sigma, \tau \in \Omega$, Then by Lemma 2.7 and Proposition 2.12, we have

$$\mu_h(J_{\sigma\tau}) \leq \eta|J_{\sigma\tau}|^h \leq \eta\xi^{3h}|J_\sigma|^h|J_\tau|^h \leq \eta^3\xi^{3h}\mu_h(J_\sigma)\mu_h(J_\eta),$$

and similarly, $\mu_h(J_{\sigma\tau}) \geq \eta^{-3}\xi^{-3h}\mu_h(J_\sigma)\mu_h(J_\eta)$. Take $L = \eta^3\xi^{3h}$. As $h > 0$, $\xi > 1$ and $\eta \geq 1$, it follows that $L > 1$, and thus

$$L^{-1}\mu_h(J_\sigma)\mu_h(J_\eta) \leq \mu_h(J_{\sigma\tau}) \leq L\mu_h(J_\sigma)\mu_h(J_\eta),$$

which is the lemma. ☐

2.14. **Topological pressure:** For $q, t \in \mathbb{R}$ and $n \geq 1$, let us write

$$Z_n(q,t) = \sum_{\sigma \in \Omega_n} (\mu_h(J_\sigma))^q \, \|\varphi'_\sigma\|^t.$$

Then for $n, p \geq 1$,

$$Z_{n+p}(q,t) = \sum_{\sigma \in \Omega_n} \sum_{\tau \in \in \Omega_p} (\mu_h(J_{\sigma\tau}))^q \, \|\varphi'_{\sigma\tau}\|^t.$$

Let us first assume $q \geq 0$. Then by Lemma 2.5 and Lemma 2.13, it follows that if $t \geq 0$ then

$$Z_{n+p}(q,t) \leq L^q Z_n(q,t) Z_p(q,t),$$

and if $t < 0$ then

$$Z_{n+p}(q,t) \leq L^q \xi^{-t} Z_n(q,t) Z_p(q,t).$$

Let us now assume $q < 0$. Using the same argument, if $t \geq 0$ then

$$Z_{n+p}(q,t) \leq L^{-q} Z_n(q,t) Z_p(q,t),$$

and if $t < 0$ then

$$Z_{n+p}(q,t) \leq L^{-q} \xi^{-t} Z_n(q,t) Z_p(q,t).$$

Hence by the standard theory of sub-additive sequences, $\lim_{k\to\infty} \frac{1}{k} \log Z_k(q,t)$ exists (cf. [F1, Corollary 1.2]). Let us denote it by $P(q,t)$, i.e.,

$$P(q,t) = \lim_{k\to\infty} \frac{1}{k} \log \sum_{\sigma \in \Omega_k} (\mu_h(J_\sigma))^q \, \|\varphi'_\sigma\|^t. \tag{6}$$

The following proposition states the well-known properties of the function $P(q,t)$ (cf. [F2, P]).

Proposition 2.15. (i) $P(q,t) : \mathbb{R} \times \mathbb{R} \to \mathbb{R}$ is continuous.
(ii) $P(q,t)$ is strictly decreasing in each variable separately.
(iii) For fixed q we have $\lim_{t\to+\infty} P(q,t) = -\infty$ and $\lim_{t\to-\infty} P(q,t) = +\infty$.
(iv) $P(q,t)$ is convex: if $q_1, q_2, t_1, t_2 \in \mathbb{R}$, $a_1, a_2 \geq 0, a_1 + a_2 = 1$, then

$$P(a_1q_1 + a_2q_2, a_1t_1 + a_2t_2) \leq a_1P(q_1,t_1) + a_2P(q_2,t_2).$$

Now for fixed q, $P(q,t)$ is a continuous function of t. Its values range from $-\infty$ (when $t \to +\infty$) to $+\infty$ (when $t \to -\infty$). Therefore, by the intermediate value theorem there is a real number β such that $P(q,\beta) = 0$. The solution β is unique, since $P(q,\cdot)$ is strictly decreasing. This defines β implicitly as a function of q: for each q there is a unique $\beta = \beta(q)$ such that $P(q,\beta(q)) = 0$.

The following proposition gives the well-known properties of the function $\beta(q)$ (cf. [F2, P]).

Proposition 2.16. Let $\beta = \beta(q)$ be defined by $P(q,\beta(q)) = 0$. Then
(i) β is a continuous function of the real variable q.
(ii) β is strictly decreasing: if $q_1 < q_2$, then $\beta(q_1) > \beta(q_2)$.
(iii) $\lim_{q\to-\infty} \beta(q) = +\infty$ and $\lim_{q\to+\infty} \beta(q) = -\infty$.
(iv) β is convex: if $q_1, q_2, a_1, a_2 \in \mathbb{R}$ with $a_1, a_2 \geq 0$ and $a_1 + a_2 = 1$, then

$$\beta(a_1q_1 + a_2q_2) \leq a_1\beta(q_1) + a_2\beta(q_2).$$

The function $\beta(q)$ is sometimes denoted by $T(q)$ and called the *temperature function.* A more general discussion of this function can be found in [HJKP], where our $\beta(q)$ function corresponds to $-\tau(q)$ in their notation.



FIGURE 1. To determine D_r first find the point of intersection of $y = \beta(q)$ and the line $y = rq$. Then D_r is the y-intercept of the line through this point and the point (1, 0).

Remark 2.17. If $q = 0$, then $P(0, \beta(0)) = 0$, which implies

$$\lim_{k\to\infty} \frac{1}{k} \log \sum_{\sigma\in\Omega_k} \|\varphi'_\sigma\|^{\beta(0)} = 0,$$

i.e., $\beta(0)$ gives the Hausdorff dimension $\dim_{\mathrm{H}}(E)$ of the cookie-cutter set E (cf. [F1]). Again

$$P(1, 0) = \lim_{k\to\infty} \frac{1}{k} \log \sum_{\sigma\in\Omega_k} \mu_h(J_\sigma) = \lim_{k\to\infty} \frac{1}{k} \log 1 = 0,$$

and hence $\beta(1) = 0$ (see Figure 1).

3. MAIN RESULT

The relationship between the quantization dimension function and the temperature function $\beta(q)$ for the Gibbs-like measure μ_h, where the temperature function is the Legendre transform of the $f(\alpha)$ curve (for the definitions of $f(\alpha)$ and the Legendre transform see [F1]) is given by the following theorem which constitutes the main result of our paper. For its graphical description see Figure 1.

Theorem 3.1. *Let μ_h be the Gibbs-like measure supported by the cookie-cutter set E. For each $r \in (0, +\infty)$ the quantization dimension (of order r) for the probability measure μ_h is given by*

$$D_r(\mu_h) = \frac{\beta(q_r)}{1 - q_r},$$

where, we recall β is the temperature function.

Lemma 3.2. *Let $0 < r < +\infty$ be fixed. Then there exists exactly one number $\kappa_r \in (0, +\infty)$ such that*

$$\lim_{k\to\infty} \frac{1}{k} \log \sum_{\sigma\in\Omega_k} (\mu_h(J_\sigma)\|\varphi'_\sigma\|^r)^{\frac{\kappa_r}{r+\kappa_r}} = 0.$$

Proof. By (6) we have

$$P(t, rt) = \lim_{k\to\infty} \frac{1}{k} \log \sum_{\sigma\in\Omega_k} \left(\mu_h(J_\sigma)\|\varphi'_\sigma\|^r\right)^t.$$

If $t = 0$, then

$$P(0, 0) = \lim_{k\to\infty} \frac{1}{k} \log \sum_{\sigma\in\Omega_k} 1 \geq \lim_{k\to\infty} \frac{1}{k} \log N^k = \log N \geq \log 2 > 0,$$

and if $t = 1$,

$$\begin{aligned} P(1, r1) &= \lim_{k\to\infty} \frac{1}{k} \log \sum_{\sigma\in\Omega_k} \mu_h(J_\sigma)\|\varphi'_\sigma\|^r \leq \lim_{k\to\infty} \frac{1}{k} \log \sum_{\sigma\in\Omega_k} \mu_h(J_\sigma) b^{-kr} \\ &= \lim_{k\to\infty} \frac{1}{k} \log \sum_{\sigma\in\Omega_k} \mu_h(J_\sigma) - r\log b = -r\log b < 0. \end{aligned}$$

Since $P(t, rt)$ is continuous and strictly decreasing, the unique $t \in \mathbb{R}$ for which $P(t, rt) = 0$ must lie between 0 and 1. Then $\kappa_r = \frac{rt}{1-t}$ satisfies the conclusion of the lemma. ☐

Lemma 3.3. *Let $0 < r < +\infty$ and κ_r be as in Lemma 3.2. Then for $n \geq 1$, we have*

$$(L\xi^r)^{-\frac{\kappa_r}{r+\kappa_r}} \leq \sum_{\sigma\in\Omega_n} \left(\mu_h(J_\sigma)\|\varphi'_\sigma\|^r\right)^{\frac{\kappa_r}{r+\kappa_r}} \leq (L\xi^r)^{\frac{\kappa_r}{r+\kappa_r}}.$$

Proof. For $\sigma \in \Omega$ let us write $s_\sigma = \mu_h(J_\sigma)\|\varphi'_\sigma\|^r$. Then for $\sigma \in \Omega_n$ and $\tau \in \Omega_p$ with $n, p \geq 1$, by Lemma 2.5 and Lemma 2.13, we have $L^{-1}\xi^{-r}s_\sigma s_\tau \leq s_{\sigma\tau} \leq L s_\sigma s_\tau \leq L\xi^r s_\sigma s_\tau$. Since $r > 0$, $L, \xi > 1$, it is true that $L^{-2}\xi^{-2r}s_\sigma s_\tau \leq s_{\sigma\tau} \leq L^2\xi^{2r}s_\sigma s_\tau$. Then by the standard theory of sub-additive sequences, $\lim_{n\to\infty} n^{-1}\log\sum_{\sigma\in\Omega_n} s_\sigma^t$ exists for any $t \in \mathbb{R}$. Let us denote this limit by $h(t)$. Hence for $t \geq 0$, we have

$$h(t) = \lim_{p\to\infty} \frac{1}{np} \log \sum_{\sigma\in\Omega_{np}} s_\sigma^t,$$

and so

$$\lim_{p\to\infty} \frac{1}{np} \log \left((L\xi^r)^{-t} \sum_{\sigma\in\Omega_n} s_\sigma^t \right)^p \leq h(t) \leq \lim_{p\to\infty} \frac{1}{np} \log \left((L\xi^r)^{t} \sum_{\sigma\in\Omega_n} s_\sigma^t \right)^p,$$

which implies

$$\frac{1}{n} \log \left((L\xi^r)^{-t} \sum_{\sigma\in\Omega_n} s_\sigma^t \right) \leq h(t) \leq \frac{1}{n} \log \left((L\xi^r)^{t} \sum_{\sigma\in\Omega_n} s_\sigma^t \right)$$

and therefore,

$$(L\xi^r)^{-t} e^{nh(t)} \leq \sum_{\sigma\in\Omega_n} s_\sigma^t \leq (L\xi^r)^t e^{nh(t)}.$$

Now substitute $t = \frac{\kappa_r}{r+\kappa_r}$ and note that $h(t) = 0$ to obtain the assertion of the lemma. ☐

We call $\Gamma \subset \Omega$ a *finite maximal anti-chain* if Γ is a finite set of words in Ω, such that every sequence in Ω_∞ is an extension of some word in Γ, but no word of Γ is an extension of another word in Γ. By $|\Gamma|$ we denote the cardinality of Γ. Note that from the definition of Γ it follows that finite maximal anti-chain does not contain the empty word $\emptyset$ as all words are extension of $\emptyset$.

Lemma 3.4. *Let Γ be a finite maximal anti-chain and κ_r satisfies the uniqueness of Lemma 3.2. Then,*

(*a*) $L^{-1}\sum_{\sigma\in\Gamma} \mu_h(J_\sigma)\mu_h \circ \varphi_\sigma^{-1} \leq \mu_h \leq L\sum_{\sigma\in\Gamma} \mu_h(J_\sigma)\mu_h \circ \varphi_\sigma^{-1}$, *and*

(*b*) $\sum_{\sigma\in\Gamma} \left(\mu_h(J_\sigma)\|\varphi'_\sigma\|^r\right)^{\frac{\kappa_r}{r+\kappa_r}} \leq (L^2\xi^r)^{\frac{\kappa_r}{r+\kappa_r}}$.

Proof. (a) Let $M = \max\{|\sigma| : \sigma \in \Gamma\}$. Note that the Borel σ-algebra on E is generated by the set $\{J_\sigma : \sigma \in \Omega\}$ of all basic intervals. For any two basic intervals either they are disjoint or one is contained in the other. Hence, it is enough to prove that for any $\omega \in \Omega_n$ with $n \geq M$,

$$L^{-1} \sum_{\sigma \in \Gamma} \mu_h(J_\sigma) \mu_h \circ \varphi_\sigma^{-1}(J_\omega) \leq \mu_h(J_\omega) \leq L \sum_{\sigma \in \Gamma} \mu_h(J_\sigma) \mu_h \circ \varphi_\sigma^{-1}(J_\omega).$$

Let $\omega \in \Omega_n$ with $n \geq M$. Since Γ is a finite maximal anti-chain, there exists $x \in \Gamma$ such that $\omega = xy$ for some $y \in \Omega$. Then, $J_\omega = J_{xy} = \varphi_{xy}(J) = \varphi_x(\varphi_y(J)) = \varphi_x(J_y)$. Hence,

$$\begin{aligned} \sum_{\sigma \in \Gamma} \mu_h(J_\sigma) \mu_h \circ \varphi_\sigma^{-1}(J_\omega) &= \sum_{\sigma \in \Gamma} \mu_h(J_\sigma) \mu_h \circ \varphi_\sigma^{-1}(\varphi_x(J_y)) = \mu_h(J_x) \mu_h \circ \varphi_x^{-1}(\varphi_x(J_y)) \\ &= \mu_h(J_x) \mu_h(J_y), \end{aligned}$$

and so by Lemma 2.13,

$$L^{-1} \sum_{\sigma \in \Gamma} \mu_h(J_\sigma) \mu_h \circ \varphi_\sigma^{-1}(J_\omega) \leq \mu_h(J_\omega) \leq L \sum_{\sigma \in \Gamma} \mu_h(J_\sigma) \mu_h \circ \varphi_\sigma^{-1}(J_\omega),$$

to obtain the assertion.

To prove (b), let $m = \min\{|\sigma| : \sigma \in \Gamma\}$. Then for each $\sigma \in \Gamma$ there exists $\tau(\sigma) \in \Omega$ with $|\tau(\sigma)| = m$ and $\tau(\sigma) \prec \sigma$, i.e., there exists $x(\sigma) \in \Omega$ such that $\sigma = \tau(\sigma)x(\sigma)$. Note that for any $\sigma \in \Gamma$ using Lemma 2.5 and Lemma 2.13, we can write

$$\|\varphi'_\sigma\| \leq \|\varphi'_{\tau(\sigma)}\| \|\varphi'_{x(\sigma)}\| \leq \|\varphi'_{\tau(\sigma)}\| \text{ and } \mu_h(J_\sigma) \leq L \mu_h(J_{\tau(\sigma)}) \mu_h(J_{x(\sigma)}) \leq L \mu_h(J_{\tau(\sigma)}).$$

Hence using Lemma 3.3, we have

$$\begin{aligned} \sum_{\sigma \in \Gamma} (\mu_h(J_\sigma) \|\varphi'_\sigma\|^r)^{\frac{\kappa_r}{r+\kappa_r}} &\leq L^{\frac{\kappa_r}{r+\kappa_r}} \sum_{\sigma \in \Gamma} \left(\mu_h(J_{\tau(\sigma)}) \|\varphi'_{\tau(\sigma)}\|^r\right)^{\frac{\kappa_r}{r+\kappa_r}} \leq L^{\frac{\kappa_r}{r+\kappa_r}} \sum_{\tau \in \Omega_m} (\mu_h(J_\tau) \|\varphi'_\tau\|^r)^{\frac{\kappa_r}{r+\kappa_r}} \\ &\leq (L^2 \xi^r)^{\frac{\kappa_r}{r+\kappa_r}}. \end{aligned}$$

□

Lemma 3.5. *Let $x, y \in J$ and $\sigma \in \Omega$. Then*

$$\xi^{-1} \|\varphi'_\sigma\| d(x, y) \leq d(\varphi_\sigma(x), \varphi_\sigma(y)) \leq \|\varphi'_\sigma\| d(x, y),$$

where ξ is the constant of Lemma 2.2.

Proof. By mean value theorem, for any $x, y \in J$ there exist some $w \in (x, y)$ such that

$$d(\varphi_\sigma(x), \varphi_\sigma(y)) = |\varphi'_\sigma(w)| d(x, y),$$

and so, by Proposition 2.4, for any $z \in J$,

$$\xi^{-1} |\varphi'_\sigma(z)| d(x, y) \leq d(\varphi_\sigma(x), \varphi_\sigma(y)) = |\varphi'_\sigma(w)| d(x, y) \leq \|\varphi'_\sigma\| d(x, y).$$

Now take the supremum over all $z \in J$, and then

$$\xi^{-1} \|\varphi'_\sigma\| d(x, y) \leq d(\varphi_\sigma(x), \varphi_\sigma(y)) \leq \|\varphi'_\sigma\| d(x, y),$$

to obtain the assertion of the lemma. □

Lemma 3.6. *Let $\Gamma \subset \Omega$ be a finite maximal anti-chain, $n \in \mathbb{N}$ with $n \geq |\Gamma|$, and $0 < r < +\infty$. Then $V_{n,r}(\mu_h) \leq \inf\left\{L \sum_{\sigma \in \Gamma} \mu_h(J_\sigma) \|\varphi'_\sigma\|^r V_{n_\sigma, r}(\mu_h) : 1 \leq n_\sigma, \ \sum_{\sigma \in \Gamma} n_\sigma \leq n\right\}$.*

Proof. Suppose $n_\sigma \geq 1$ for each $\sigma \in \Gamma$, and $\sum_{\sigma\in\Gamma} n_\sigma \leq n$. For each $\sigma \in \Gamma$ let α_σ be an n_σ-optimal set for $V_{n_\sigma,r}(\mu_h)$. Since $|\cup_{\sigma\in\Gamma}\varphi_\sigma(\alpha_\sigma)| \leq n$ and $\mu_h \leq L\sum_{\sigma\in\Gamma}\mu_h(J_\sigma)\mu_h\circ\varphi_\sigma^{-1}$, we have

$$\begin{aligned}
V_{n,r}(\mu_h) &\leq \int d(x, \bigcup \varphi_\sigma(\alpha_\sigma))^r d\mu_h(x) \\
&\leq L\sum_{\sigma\in\Gamma}\mu_h(J_\sigma)\int d(x, \bigcup \varphi_\sigma(\alpha_\sigma))^r d(\mu_h\circ\varphi_\sigma^{-1})(x) \\
&= L\sum_{\sigma\in\Gamma}\mu_h(J_\sigma)\int d(\varphi_\sigma(x), \varphi_\sigma(\alpha_\sigma))^r d\mu_h(x) \\
&\leq L\sum_{\sigma\in\Gamma}\mu_h(J_\sigma)\|\varphi'_\sigma\|^r\int d(x, \alpha_\sigma)^r d\mu_h(x) \\
&= L\sum_{\sigma\in\Gamma}\mu_h(J_\sigma)\|\varphi'_\sigma\|^r V_{n_\sigma,r}(\mu_h),
\end{aligned}$$

which implies the lemma. □

Proposition 3.7. *Let* $0 < r < +\infty$ *and* κ_r *be as in Lemma 3.2. Then* $\limsup_{n\to\infty} nV_{n,r}^{\frac{\kappa_r}{h}}(\mu_h) < +\infty$.

Proof. Let $\epsilon_0 = \inf\{\left(\mu_h(J_j)\|\varphi'_j\|^r\right)^{\frac{\kappa_r}{r+\kappa_r}} : 1 \leq j \leq N\}$. Since

$$1 > \mu_h(J_j)\|\varphi'_j\|^r \geq \eta^{-1}|J_j|B^{-r} \geq \eta^{-1}\xi^{-2}B^{-1}B^{-r} > 0,$$

we have $0 < \epsilon_0 < 1$. Fix $m \in \mathbb{N}$. Choose any $n \in \mathbb{N}$ so that $\frac{m}{n}\left(L^3\xi^{2r}\right)^{\frac{\kappa_r}{r+\kappa_r}} < \epsilon_0^2$, and set $\epsilon = \epsilon_0^{-1}\frac{m}{n}\left(L^3\xi^{2r}\right)^{\frac{\kappa_r}{r+\kappa_r}}$. Then $0 < \epsilon < 1$. Let

$$\Gamma = \Gamma(\epsilon) = \{\sigma \in \Omega : (\mu_h(J_\sigma)\|\varphi'_\sigma\|^r)^{\frac{\kappa_r}{r+\kappa_r}} < \epsilon \leq (\mu_h(J_{\sigma^-})\|\varphi'_{\sigma^-}\|^r)^{\frac{\kappa_r}{r+\kappa_r}}\}.$$

By Lemma 3.4, we have

$$\begin{aligned}
&(L^2\xi^r)^{\frac{\kappa_r}{r+\kappa_r}} \\
&\geq \sum_{\sigma\in\Gamma}(\mu_h(J_\sigma)\|\varphi'_\sigma\|^r)^{\frac{\kappa_r}{r+\kappa_r}} \geq (L\xi^r)^{-\frac{\kappa_r}{r+\kappa_r}}\sum_{\sigma\in\Gamma}(\mu_h(J_{\sigma^-})\|\varphi'_{\sigma^-}\|^r)^{\frac{\kappa_r}{r+\kappa_r}}\left(\mu_h(J_{\sigma_{|\sigma|}})\|\varphi'_{\sigma_{|\sigma|}}\|^r\right)^{\frac{\kappa_r}{r+\kappa_r}} \\
&\geq (L\xi^r)^{-\frac{\kappa_r}{r+\kappa_r}}|\Gamma|\epsilon\epsilon_0,
\end{aligned}$$

which implies $|\Gamma| \leq (\epsilon\epsilon_0)^{-1}(L^3\xi^{2r})^{\frac{\kappa_r}{r+\kappa_r}} = \frac{n}{m} < \infty$, i.e., Γ is a finite maximal anti-chain and $n \geq |\Gamma|m$. Hence by the previous lemma, we have

$$\begin{aligned}
V_{n,r}(\mu_h) &\leq L\sum_{\sigma\in\Gamma}\mu_h(J_\sigma)\|\varphi'_\sigma\|^r V_{m,r}(\mu_h) \\
&= \sum_{\sigma\in\Gamma}(\mu_h(J_\sigma)\|\varphi'_\sigma\|^r)^{\frac{\kappa_r}{r+\kappa_r}}(\mu_h(J_\sigma)\|\varphi'_\sigma\|^r)^{\frac{r}{r+\kappa_r}}V_{m,r}(\mu_h) \\
&< \sum_{\sigma\in\Gamma}(\mu_h(J_\sigma)\|\varphi'_\sigma\|^r)^{\frac{\kappa_r}{r+\kappa_r}}\epsilon^{\frac{r}{\kappa_r}}V_{m,r}(\mu_h) \\
&\leq (L^2\xi^r)^{\frac{\kappa_r}{r+\kappa_r}}\epsilon^{\frac{r}{\kappa_r}}V_{m,r}(\mu_h) \\
&= (L^2\xi^r)^{\frac{\kappa_r}{r+\kappa_r}}\epsilon_0^{-\frac{r}{\kappa_r}}\left(\frac{m}{n}\right)^{\frac{r}{\kappa_r}}\left(L^3\xi^{2r}\right)^{\frac{r}{r+\kappa_r}}V_{m,r}(\mu_h),
\end{aligned}$$

which implies $nV_{n,r}^{\frac{\kappa_r}{r}}(\mu_h) \leq (L^2\xi^r)^{\frac{\kappa_r^2}{r(r+\kappa_r)}} (L^3\xi^{2r})^{\frac{\kappa_r}{r+\kappa_r}} \epsilon_0^{-1} m V_{m,r}^{\frac{\kappa_r}{r}}(\mu_h)$. Since for fixed m, this inequality holds for all but a finite number of n, we have

$$\limsup_{n\to\infty} nV_{n,r}^{\frac{\kappa_r}{r}}(\mu_h) \leq (L^2\xi^r)^{\frac{\kappa_r^2}{r(r+\kappa_r)}} \left(L^3\xi^{2r}\right)^{\frac{\kappa_r}{r+\kappa_r}} \epsilon_0^{-1} m V_{m,r}^{\frac{\kappa_r}{r}}(\mu_h) < +\infty.$$

$\square$

Note 3.8. We say that the cookie-cutter mapping f satisfies the *open set condition* (OSC) if the corresponding set of branch inverses $\{\varphi_1, \varphi_2, \cdots, \varphi_N\}$ satisfies the open set condition: there exists a bounded nonempty open set $U \subset J$ (in the topology of J) such that $\varphi_j(U) \subset U$, and $\varphi_i(U) \cap \varphi_j(U) = \emptyset$ for $1 \leq i \neq j \leq N$. Furthermore, f satisfies the *strong open set condition* (SOSC) if U can be chosen such that $U \cap E \neq \emptyset$. Note that we can choose $U = (0, 1)$ and so, because of the net properties of the basic intervals of the cookie-cutter set, it follows that the cookie-cutter mapping f satisfies the strong open set condition.

Lemma 3.9. *Let $\Gamma \subset \Omega$ be a finite maximal anti-chain. Then there exists $n_0 = n_0(\Gamma)$ such that for every $n \geq n_0$ there exists a set of positive integers $\{n_\sigma := n_\sigma(n)\}_{\sigma\in\Gamma}$ such that $\sum_{\sigma\in\Gamma} n_\sigma \leq n$ and*

$$u_{n,r}(\mu_h) \geq (L\xi^r)^{-1} \sum_{\sigma\in\Gamma} \mu_h(J_\sigma)\|\varphi'_\sigma\|^r u_{n_\sigma,r}(\mu_h).$$

Proof. Let U be the open set from the strong open set condition. Then there exists $\tau \in \Omega$ such that $\varphi_\tau(J) \subset U$. Let $\epsilon = d(\varphi_\tau(J), U^c)$ and $\delta = \min\{\xi^{-1}\|\varphi'_\sigma\| : \sigma \in \Gamma\}$. Then for $\sigma \in \Gamma$ we have $d(\varphi_\sigma\varphi_\tau(J), \varphi_\sigma(U^c)) \geq \xi^{-1}\|\varphi'_\sigma\| d(\varphi_\tau(J), U^c) \geq \delta\epsilon$, which implies $d(x, U^c) \geq d(x, \varphi_\sigma(U^c)) \geq \delta\epsilon$ for any $x \in \varphi_\sigma(\varphi_\tau(J))$. For each n, let α_n be an n-optimal set for $u_{n,r}$ and let $\delta_n = \max\{d(x, \alpha_n \cup U^c) : x \in E\}$. Since $\delta_n \to 0$ as $n \to \infty$ we can choose n_0 such that $\delta_n < \delta\epsilon$ for all $n \geq n_0$. Suppose $n \geq n_0$ and $x \in \varphi_\sigma(\varphi_\tau(E))$. Note that $\varphi_\sigma(\varphi_\tau(E)) \subset \varphi_\tau(E) \subset \varphi_\tau(J) \subset U$. Then there exists $a \in \alpha_n \cup U^c$ such that $d(x, \alpha_n \cup U^c) = d(x, a) \leq \delta_n < \delta\epsilon$, and so $a \notin \varphi_\sigma(U^c)$, i.e., $a \in \varphi_\sigma(U)$. Therefore, letting $\alpha_{n_\sigma} = \alpha_n \cap \varphi_\sigma(U)$, we have $n_\sigma := |\alpha_{n_\sigma}| \geq 1$ and $\sum_{\sigma\in\Gamma} n_\sigma \leq n$. For any $x \in E$, we claim that there exists $y \in \alpha_{n_\sigma} \cup \varphi_\sigma(U^c)$ such that $d(\varphi_\sigma(x), \alpha_{n_\sigma} \cup \varphi_\sigma(U^c)) = d(x, y)$. If not let $y \notin \alpha_{n_\sigma} \cup \varphi_\sigma(U^c)$. Then $y \notin \alpha_n \cap \varphi_\sigma(U)$ and $y \notin \varphi_\sigma(U^c)$, i.e., $z := \varphi_\sigma^{-1}(y) \in J$ be such that $z \notin U$ and $z \notin U^c$, which gives a contradiction and thus the claim is true, and it implies $d(\varphi_\sigma(x), \alpha_n \cup \varphi_\sigma(U^c)) = d(\varphi_\sigma(x), \alpha_{n_\sigma} \cup \varphi_\sigma(U^c))$. Again for any $x \in E$ as $U^c \subset (\varphi_\sigma(U))^c = \varphi_\sigma(U^c)$, we have $d(\varphi_\sigma(x), \alpha_n \cup U^c) \geq d(\varphi_\sigma(x), \alpha_n \cup \varphi_\sigma(U^c))$. Hence,

$$\begin{aligned} u_{n,r}(\mu_h) &= \int d(x, \alpha_n \cup U^c)^r d\mu_h(x) \geq L^{-1} \sum_{\sigma\in\Gamma} \mu_h(J_\sigma) \int d(\varphi_\sigma(x), \alpha_n \cup U^c)^r d\mu_h(x) \\ &\geq L^{-1} \sum_{\sigma\in\Gamma} \mu_h(J_\sigma) \int d(\varphi_\sigma(x), \alpha_n \cup \varphi_\sigma(U^c))^r d\mu_h(x) \\ &= L^{-1} \sum_{\sigma\in\Gamma} \mu_h(J_\sigma) \int d(\varphi_\sigma(x), \alpha_{n_\sigma} \cup \varphi_\sigma(U^c))^r d\mu_h(x) \\ &\geq (L\xi^r)^{-1} \sum_{\sigma\in\Gamma} \mu_h(J_\sigma)\|\varphi'_\sigma\|^r \int d(x, \varphi_\sigma^{-1}(\alpha_{n_\sigma}) \cup U^c)^r d\mu_h(x) \\ &\geq (L\xi^r)^{-1} \sum_{\sigma\in\Gamma} \mu_h(J_\sigma)\|\varphi'_\sigma\|^r u_{n_\sigma,r}(\mu_h). \end{aligned}$$

$\square$

Proposition 3.10. *Let μ_h be the Gibbs-like measure, and let $0 < r < +\infty$. Moreover, let κ_r be as in Lemma 3.2, and $0 < \ell < \kappa_r$. Then $\liminf_{n\to\infty} nV_{n,r}^{\frac{\ell}{r}}(\mu_h) > 0$.*

Proof. Since $0 < \ell < \kappa_r$ and κ_r is unique for which

$$\lim_{k\to\infty} \frac{1}{k} \log \sum_{\sigma\in\Omega_k} (\mu_h(J_\sigma)\|\varphi'_\sigma\|^r)^{\frac{\kappa_r}{r+\kappa_r}} = 0,$$

we have

$$\sum_{\sigma\in\Omega_m} (\mu_h(J_\sigma)\|\varphi'_\sigma\|^r)^{\frac{\ell}{r+\ell}} \to \infty \text{ as } m \to \infty.$$

Choose m so that the above sum is greater than $L\xi^r$ and let $\Gamma = \{\sigma \in \Omega : |\sigma| = m\}$. Then Γ is a finite maximal anti-chain. By Lemma 3.9, we have n_0 and for $n \geq n_0$ the numbers $\{n_\sigma := n_\sigma(n)\}_{\sigma\in\Gamma}$ which satisfy the conclusion of the lemma. Set $c = \min\{n^{r/\ell}u_{n,r}(\mu_h) : n \leq n_0\}$. Clearly each $u_{n,r}(\mu_h) > 0$ and hence $c > 0$. Suppose $n \geq n_0$ and $k^{r/\ell}u_{k,r}(\mu_h) \geq c$ for all $k < n$. Hence using Lemma 3.9, we have

$$\begin{aligned} n^{r/\ell}u_{n,r}(\mu_h) &\geq n^{r/\ell}(L\xi^r)^{-1} \sum_{\sigma\in\Gamma} \mu_h(J_\sigma)\|\varphi'_\sigma\|^r u_{n_\sigma,r}(\mu_h) \\ &= n^{r/\ell}(L\xi^r)^{-1} \sum_{\sigma\in\Gamma} \mu_h(J_\sigma)\|\varphi'_\sigma\|^r (n_\sigma(n))^{-r/\ell}(n_\sigma(n))^{r/\ell}u_{n_\sigma,r}(\mu_h) \\ &\geq c(L\xi^r)^{-1} \sum_{\sigma\in\Gamma} \mu_h(J_\sigma)\|\varphi'_\sigma\|^r \left(\frac{n_\sigma(n)}{n}\right)^{-r/\ell}. \end{aligned}$$

Using Hölder's inequality (with exponents less than 1), we have

$$n^{r/\ell}u_{n,r}(\mu_h) \geq c(L\xi^r)^{-1} \left(\sum_{\sigma\in\Gamma} (\mu_h(J_\sigma)\|\varphi'_\sigma\|^r)^{\ell/(r+\ell)}\right)^{(1+r/\ell)} \left(\sum_{\sigma\in\Gamma} (\frac{n_\sigma(n)}{n})^{(-r/\ell)(-\ell/r)}\right)^{-r/\ell}.$$

By our choice of Γ, which depends only on ℓ and not on n, and the fact that $\sum_{\omega\in\Gamma} n_\omega(n) \leq n$, we have $n^{r/\ell}u_{n,r}(\mu_h) \geq c$. Therefore, by induction,

$$\liminf_{n\to\infty} n u_{n,r}^{\ell/r}(\mu_h) \geq c^{\ell/r} > 0, \text{ i.e., } \liminf n V_{n,r}^{\frac{\ell}{r}}(\mu_h) > 0,$$

and thus the lemma is yielded. □

Proof of Theorem 3.1. Note that $e_{n,r} = V_{n,r}^{\frac{1}{r}}$, and by [GL1, Proposition 11.3]) we know:

(a) If $0 \leq t < \underline{D}_r < s$ then

$$\lim_{n\to\infty} n e_{n,r}^t = +\infty \text{ and } \liminf_{n\to\infty} n e_{n,r}^s = 0.$$

(b) If $0 \leq t < \overline{D}_r < s$ then

$$\limsup_{n\to\infty} n e_{n,r}^t = +\infty \text{ and } \lim_{n\to\infty} n e_{n,r}^s = 0.$$

From (a) and Proposition 3.10, we have $\ell \leq \underline{D}_r$ whenever $\ell < \kappa_r$. Hence, $\kappa_r \leq \underline{D}_r$. From ($b$) and Proposition 3.7, we have $\overline{D}_r \leq \kappa_r$. Hence, $\kappa_r \leq \underline{D}_r \leq \overline{D}_r \leq \kappa_r$, i.e., the quantization dimension D_r exists and $D_r = \kappa_r$. Note that if $q_r = \frac{\kappa_r}{r+\kappa_r}$, by Lemma 3.2, we have $\beta(q_r) = rq_r$, and then $D_r = \frac{\beta(q_r)}{1-q_r}$. Thus the proof of the theorem is complete.

References

[BW] J.A. Bucklew and G.L. Wise, *Multidimensional asymptotic quantization with rth power distortion measures*, IEEE Trans. Inform. Theory 28, 239-247 (1982).

[F1] K.J. Falconer, *Techniques in fractal geometry*, Chichester: Wiley, 1997.

[F2] K.J. Falconer, *The multifractal spectrum of statistically self-similar measures*, Journal of Theoretical Probability, Vol 7, No. 3, 681-701, 1994.

[GG] A. Gersho and R.M. Gray, *Vector quantization and signal compression*, Kluwer Academy publishers: Boston, 1992.

[GN] R. Gray and D. Neuhoff, *Quantization,* IEEE Trans. Inform. Theory 44, 2325-2383 (1998).

[GL1] S. Graf and H. Luschgy, *Foundations of quantization for probability distributions*, Lecture Notes in Mathematics 1730, Springer, Berlin, 2000.

[GL2] S. Graf and H. Luschgy, *The Quantization dimension of self-similar probabilities*, Math. Nachr., 241 (2002), 103-109.

[GL3] S. Graf and H. Luschgy, *Asymptotics of the Quantization Errors for Self-Similar Probabilities,* Real Anal. Exchange, Volume 26, Number 2 (2000), 795-810.

[H] J. Hutchinson, *Fractals and self-similarity*, Indiana Univ. J., 30, 713-747, 1981.

[HJKP] T. Halsey, M. Jensen, L. Kadanoff and I. Procaccia, *Fractal measures and their singularities: the characterization of strange sets*, Phys. Ref. A, A 33(2) 1141-51, 1986.

[LM] L.J. Lindsay and R.D. Mauldin, *Quantization dimension for conformal iterated function systems*, Nonlinearity, 15 (2002), 189-199.

[P] N. Patzschke, *Self-conformal multifractal measures*, Adv. Appl. Math., Volume 19, Issue 4 (1997), 486-513.

[R1] M.K. Roychowdhury, *Quantization dimension for some Moran measures*, Proc. Amer. Math. Soc., 138 (2010), 4045-4057.

[R2] M.K. Roychowdhury, *Quantization dimension function and Gibbs measure associated with Moran set*, J. Math. Anal. Appl., 373 (2011), 73-82.

[WD] X. Wang and M. Dai, *Mixed Quantization Dimension Function and Temperature Function for Conformal Measures*, International Journal of Nonlinear Science, Vol.10(2010), 24-31.

[Z] P.L. Zador, *Development and evaluation of procedures for quantizing multivariate distributions,* PhD Thesis, Stanford University (1964).

[Y] K. Yoshida, *Functional Analysis*, Berlin-Heidelberg-New York: Springer. 1966.

Department of Mathematics, The University of Texas-Pan American, 1201 West University Drive, Edinburg, TX 78539-2999, USA.

E-mail address: `roychowdhurymk@utpa.edu`